arXiv:0811.2091v1 [math.FA] 13 Nov 2008

GROWTH ESTIMATES FOR A CLASS OF SUBHARMONIC FUNCTIONS IN A HALF SPACE *

PAN GUOSHUANG[1,2] AND DENG GUANTIE[1,**]

Abstract. A class of subharmonic functions represented by the modified kernels are proved to have the growth estimates $u(x) = o(x_n^{1-\alpha}|x|^{m+\alpha})$ at infinity in the upper half space of $\mathbf{R}^n$, which generalizes the growth properties of analytic functions and harmonic functions.

1. Introduction and Main Theorem

Let $\mathbf{R}^n (n \geq 3)$ denote the n-dimensional Euclidean space with points $x = (x_1, x_2, \cdots, x_{n-1}, x_n) = (x', x_n)$, where $x' \in \mathbf{R}^{n-1}$ and $x_n \in \mathbf{R}$. The boundary and closure of an open Ω of $\mathbf{R}^n$ are denoted by $\partial\Omega$ and $\overline{\Omega}$ respectively. The upper half-space H is the set $H = \{x = (x', x_n) \in \mathbf{R}^n : x_n > 0\}$, whose boundary is ∂H . We write $B(x, \rho)$ and $\partial B(x, \rho)$ for the open ball and the sphere of radius ρ centered at x in $\mathbf{R}^n$. We identify $\mathbf{R}^n$ with $\mathbf{R}^{n-1} \times \mathbf{R}$ and $\mathbf{R}^{n-1}$ with $\mathbf{R}^{n-1} \times \{0\}$, with this convention we then have $\partial H = \mathbf{R}^{n-1}$, writing typical points $x,\ y \in \mathbf{R}^n$ as $x = (x', x_n),\ y = (y', y_n)$, where $x' = (x_1, x_2, \cdots, x_{n-1}),\ y' = (y_1, y_2, \cdots y_{n-1}) \in \mathbf{R}^{n-1}$ and putting

$$x \cdot y = \sum_{j=1}^{n} x_j y_j = x' \cdot y' + x_n y_n, \quad |x| = \sqrt{x \cdot x}, \quad |x'| = \sqrt{x' \cdot x'}.$$

For $x \in \mathbf{R}^n \backslash \{0\}$, let([**16**])

$$E(x) = -r_n |x|^{2-n},$$

where $|x|$ is the Euclidean norm, $r_n = \frac{1}{(n-2)\omega_n}$ and $\omega_n = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$ is the surface area of the unit sphere in $\mathbf{R}^n$. We know that E is locally integrable in $\mathbf{R}^n$.

Key words and phrases. Subharmonic function, Modified Poisson kernel, Modified Green function, Growth estimate.

2000 Mathematics Subject Classification. 31B05, 31B10.

$*$ The project supported by NSFC (Grant No.10671022) and by RFDP (Grant No.20060027023)

$**$:Corresponding author.

The Green function $G(x,y)$ for the upper half space H is given by([**16**])

$$G(x,y)=E(x-y)-E(x-y^*) \qquad x,y\in\overline{H},\ x\neq y,$$

where * denotes reflection in the boundary plane ∂H just as $y^*=(y_1,y_2,\cdots,y_{n-1},-y_n)$, then we define the Poisson kernel $P(x,y')$ when $x\in H$ and $y'\in\partial H$ by

$$P(x,y')=-\frac{\partial G(x,y)}{\partial y_n}\bigg|_{y_n=0}=\frac{2x_n}{\omega_n|x-(y',0)|^n}.$$

The Dirichlet problem of upper half space is to find a function u satisfying

$$u\in C^2(H), \tag{1.1}$$

$$\Delta u=0, x\in H, \tag{1.2}$$

$$\lim_{x\to x'}u(x)=f(x') \text{ nontangentially a.e.} x'\in\partial H, \tag{1.3}$$

where f is a measurable function of $\mathbf{R}^{n-1}$. The Poisson integral of the upper half space is defined by

$$u(x)=P[f](x)=\int_{\mathbf{R}^{n-1}}P(x,y')f(y')dy'.$$

As we all know, the Poisson integral $P[f]$ exists if

$$\int_{\mathbf{R}^{n-1}}\frac{|f(y')|}{1+|y'|^n}dy'<\infty.$$

(see [**17,18**] and [**20**])In this paper, we will consider measurable functions f in $\mathbf{R}^{n-1}$ satisfying

$$\int_{\mathbf{R}^{n-1}}\frac{|f(y')|}{1+|y'|^{n+m}}dy'<\infty. \tag{1.4}$$

It is well known that the Poisson kernel $P(x,y')$ has a series expansion in terms of the ultraspherical (or Gegenbauer) polynomials $C_k^\lambda(t)$ $(\lambda=\frac{n}{2})$([**7**] and [**12**]). The latter can be defined by a generating function

$$(1-2tr+r^2)^{-\lambda}=\sum_{k=0}^{\infty}C_k^\lambda(t)r^k, \tag{1.5}$$

where $|r|<1$, $|t|\leq 1$ and $\lambda>0$. The coefficients $C_k^\lambda(t)$ is called the ultraspherical (or Gegenbauer) polynomial of degree k associated with λ, the function $C_k^\lambda(t)$ is a polynomial of degree k in t. To obtain a solution of Dirichlet problem for the boundary date f, as in [**3,5,15**] and [**20**], we use the following modified functions defined by

$$E_m(x-y)=\begin{cases} E(x-y) & \text{when } |y|\leq 1,\\ E(x-y)+\sum_{k=0}^{m-1}\frac{r_n|x|^k}{|y|^{n-2+k}}C_k^{\frac{n-2}{2}}\left(\frac{x\cdot y}{|x||y|}\right) & \text{when } |y|>1.\end{cases}$$

Then we can define modified Green function $G_m(x,y)$ and the modified Poisson kernel $P_m(x,y')$ by([**1,2,4,11**] and [**20**])

$$G_m(x,y) = E_{m+1}(x-y) - E_{m+1}(x-y^*) \qquad x,y \in \overline{H},\ x \neq y; \quad (1.6)$$

$$P_m(x,y') = \begin{cases} P(x,y') & \text{when } |y'| \leq 1, \\ P(x,y') - \sum_{k=0}^{m-1} \frac{2x_n|x|^k}{\omega_n|y'|^{n+k}} C_k^{n/2}\left(\frac{x\cdot(y',0)}{|x||y'|}\right) & \text{when } |y'| > 1. \end{cases} \quad (1.7)$$

Siegel-Talvila([**3**]) have proved the following result:

Theorem A Let f be a measurable function in $\mathbf{R}^{n-1}$ satisfying (1.4), then the harmonic function

$$v(x) = \int_{\mathbf{R}^{n-1}} P_m(x,y')f(y')dy' \quad x \in H \quad (1.8)$$

satisfies (1.1), (1.2), (1.3) and

$$v(x) = o(x_n^{1-n}|x|^{m+n}) \quad \text{as } |x| \to \infty. \quad (1.9)$$

where $P_m(x,y')$ is defined by (1.7).

In order to describe the asymptotic behaviour of subharmonic functions in half-spaces([**8,9**] and [**10**]), we establish the following theorems.

Theorem 1 Let f be a measurable function in $\mathbf{R}^{n-1}$ satisfying (1.4), and $0 < \alpha \leq n$. Let $v(x)$ be the harmonic function defined by (1.8). Then there exists $x_j \in H,\ \rho_j > 0$, such that

$$\sum_{j=1}^{\infty} \frac{\rho_j^{n-\alpha}}{|x_j|^{n-\alpha}} < \infty \quad (1.10)$$

holds and

$$v(x) = o(x_n^{1-\alpha}|x|^{m+\alpha}) \quad \text{as } |x| \to \infty \quad (1.11)$$

holds in $H - G$. where $G = \bigcup_{j=1}^{\infty} B(x_j, \rho_j)$.

Remark If $\alpha = n$, then (1.10) is a finite sum, the set G is the union of finite balls, so (1.9) holds in H. This is just the result of Siegel-Talvila, therefore, our result (1.11) is the generalization of Theorem A.

Next, we will generalize Theorem 1 to subharmonic functions.

Theorem 2 Let f be a measurable function in $\mathbf{R}^{n-1}$ satisfying (1.4), let μ be a positive Borel measure satisfying

$$\int_H \frac{y_n}{1+|y|^{n+m}} d\mu(y) < \infty. \quad (1.12)$$

Write the subharmonic function

$$u(x) = v(x) + h(x), \quad x \in H$$

where $v(x)$ is the harmonic function defined by (1.8), $h(x)$ is defined by

$$h(x) = \int_H G_m(x,y)d\mu(y)$$

and $G_m(x,y)$ is defined by (1.6). Then there exists $x_j \in H$, $\rho_j > 0$, such that (1.10) holds and

$$u(x) = o(x_n^{1-\alpha}|x|^{m+\alpha}) \quad \text{as } |x| \to \infty$$

holds in $H - G$, where $G = \bigcup_{j=1}^{\infty} B(x_j, \rho_j)$ and $0 < \alpha < 2$.

Next we are concerned with minimal thinness at infinity for $v(x)$ and $h(x)$, for a set $E \subset H$ and an open set $F \subset \mathbf{R}^{n-1}$, we consider the capacity

$$C(E;F) = \inf \int_{\mathbf{R}^{n-1}} g(y')dy'$$

where the infimum is taken over all nonnegative measurable functions g such that $g = 0$ outside F and

$$\int_{\mathbf{R}^{n-1}} \frac{g(y')}{|x-(y',0)|^n} dy' \geq 1 \qquad \text{for all } x \in E.$$

We say that $E \subset H$ is minimally thin at infinity if

$$\sum_{i=1}^{\infty} 2^{-in} C(E_i;F_i) < \infty,$$

where $E_i = \{x \in E : 2^i \leq |x| < 2^{i+1}\}$ and $F_i = \{x \in \mathbf{R}^{n-1} : 2^i < |x| < 2^{i+3}\}$.

Theorem 3 Let f be a measurable function in $\mathbf{R}^{n-1}$ satisfying (1.4), then there exists a set $E \subset H$ such that E is minimally thin at infinity and

$$\lim_{|x|\to\infty, x \in H-E} \frac{v(x)}{x_n|x|^m} = 0.$$

Similarly, for $h(x)$, we can also conclude the following:

Corollary 1 Let μ be a positive Borel measure satisfying (1.12), then there exists a set $E \subset H$ such that E is minimally thin at infinity and

$$\lim_{|x|\to\infty, x \in H-E} \frac{h(x)}{x_n|x|^m} = 0.$$

Finally we are concerned with rarefiedness at infinity for $v(x)$ and $h(x)$, for a set $E \subset H$ and an open set $F \subset H$, we consider the capacity

$$C(E;F) = \inf \int_H g(y)d\mu(y)$$

where the infimum is taken over all nonnegative measurable functions g such that $g = 0$ outside F and

$$\int_H \frac{g(y)}{|x-y|^{n-1}} d\mu(y) \geq 1 \qquad \text{for all } x \in E.$$

We say that $E \subset H$ is rarefied at infinity if

$$\sum_{i=1}^{\infty} 2^{-i(n-1)} C(E_i; F_i) < \infty,$$

where E_i is as in Theorem 3 and $F_i = \{x \in H : 2^i < |x| < 2^{i+3}\}$.

Theorem 4 Let μ be a positive Borel measure satisfying (1.12), then there exists a set $E \subset H$ such that E is rarefied at infinity and

$$\lim_{|x|\to\infty, x\in H-E} \frac{h(x)}{|x|^{m+1}} = 0.$$

Similarly, for $v(x)$, we can also conclude the following:

Corollary 2 Let f be a measurable function in $\mathbf{R}^{n-1}$ satisfying (1.4), then there exists a set $E \subset H$ such that E is rarefied at infinity and

$$\lim_{|x|\to\infty, x\in H-E} \frac{v(x)}{|x|^{m+1}} = 0.$$

2. PROOF OF THEOREM

Let μ be a positive Borel measure in $\mathbf{R}^n$, $\beta \geq 0$, the maximal function $M(d\mu)(x)$ of order β is defined by

$$M(d\mu)(x) = \sup_{0<r<\infty} \frac{\mu(B(x,r))}{r^\beta},$$

then the maximal function $M(d\mu)(x) : \mathbf{R}^n \to [0, \infty)$ is lower semicontinuous, hence measurable. To see this, for any $\lambda > 0$, let $D(\lambda) = \{x \in \mathbf{R}^n : M(d\mu)(x) > \lambda\}$. Fix $x \in D(\lambda)$, then there exists $r > 0$ such that $\mu(B(x,r)) > tr^\beta$ for some $t > \lambda$, and there exists $\delta > 0$ satisfying $(r+\delta)^\beta < \frac{tr^\beta}{\lambda}$. If $|y - x| < \delta$, then $B(y, r+\delta) \supset B(x,r)$, therefore $\mu(B(y, r+\delta)) \geq tr^\beta > \lambda(r+\delta)^\beta$. Thus $B(x,\delta) \subset D(\lambda)$. This proves that $D(\lambda)$ is open for each $\lambda > 0$.

In order to obtain the results, we need these lemmas below:

Lemma 1 Let μ be a positive Borel measure in $\mathbf{R}^n$, $\beta \geq 0$, $\mu(\mathbf{R}^n) < \infty$, for any $\lambda \geq 5^\beta \mu(\mathbf{R}^n)$, set

$$E(\lambda) = \{x \in \mathbf{R}^n : |x| \geq 2, M(d\mu)(x) > \frac{\lambda}{|x|^\beta}\}$$

then there exists $x_j \in E(\lambda)$, $\rho_j > 0$, $j = 1, 2, \cdots$, such that

$$E(\lambda) \subset \bigcup_{j=1}^{\infty} B(x_j, \rho_j) \tag{2.1}$$

and

$$\sum_{j=1}^{\infty}\frac{\rho_j^{\beta}}{|x_j|^{\beta}}\leq\frac{3\mu(\mathbf{R}^n)5^{\beta}}{\lambda}. \tag{2.2}$$

Proof: Let $E_k(\lambda)=\{x\in E(\lambda):2^k\leq|x|<2^{k+1}\}$, then for any $x\in E_k(\lambda)$, there exists $r(x)>0$, such that $\mu(B(x,r(x)))>\lambda(\frac{r(x)}{|x|})^{\beta}$, therefore $r(x)\leq 2^{k-1}$. Since $E_k(\lambda)$ can be covered by the union of a family of balls $\{B(x,r(x)):x\in E_k(\lambda)\}$, by the Vitali Lemma([**6**]), there exists $\Lambda_k\subset E_k(\lambda)$, Λ_k is at most countable, such that $\{B(x,r(x)):x\in\Lambda_k\}$ are disjoint and

$$E_k(\lambda)\subset\cup_{x\in\Lambda_k}B(x,5r(x)),$$

so

$$E(\lambda)=\cup_{k=1}^{\infty}E_k(\lambda)\subset\cup_{k=1}^{\infty}\cup_{x\in\Lambda_k}B(x,5r(x)).$$

On the other hand, note that $\cup_{x\in\Lambda_k}B(x,r(x))\subset\{x:2^{k-1}\leq|x|<2^{k+2}\}$, so that

$$\sum_{x\in\Lambda_k}\frac{(5r(x))^{\beta}}{|x|^{\beta}}\leq 5^{\beta}\sum_{x\in\Lambda_k}\frac{\mu(B(x,r(x)))}{\lambda}\leq\frac{5^{\beta}}{\lambda}\mu\{x:2^{k-1}\leq|x|<2^{k+2}\}.$$

Hence we obtain

$$\sum_{k=1}^{\infty}\sum_{x\in\Lambda_k}\frac{(5r(x))^{\beta}}{|x|^{\beta}}\leq\sum_{k=1}^{\infty}\frac{5^{\beta}}{\lambda}\mu\{x:2^{k-1}\leq|x|<2^{k+2}\}\leq\frac{3\mu(\mathbf{R}^n)5^{\beta}}{\lambda}.$$

Rearrange $\{x:x\in\Lambda_k,k=1,2,\cdots\}$ and $\{5r(x):x\in\Lambda_k,k=1,2,\cdots\}$, we get $\{x_j\}$ and $\{\rho_j\}$ such that (2.1) and (2.2) hold.

Lemma 2 Gegenbauer polynomials have the following properties:
(1) $|C_k^{\lambda}(t)|\leq C_k^{\lambda}(1)=\frac{\Gamma(2\lambda+k)}{\Gamma(2\lambda)\Gamma(k+1)},\quad |t|\leq 1$;
(2) $\frac{d}{dt}C_k^{\lambda}(t)=2\lambda C_{k-1}^{\lambda+1}(t),\quad k\geq 1$;
(3) $\sum_{k=0}^{\infty}C_k^{\lambda}(1)r^k=(1-r)^{-2\lambda}$;
(4) $|C_k^{\frac{n-2}{2}}(t)-C_k^{\frac{n-2}{2}}(t^*)|\leq(n-2)C_{k-1}^{n/2}(1)\,|t-t^*|,\quad |t|\leq 1,\quad |t^*|\leq 1$.
Proof: (1) and (2) can be derived from [**7**] and [**13**]; (3) follows by taking $t=1$ in (1.5); (4) follows by (1), (2) and the Mean Value Theorem for Derivatives.

Lemma 3 Green function $G(x,y)$ has the following estimates:
(1) $|G(x,y)|\leq\frac{r_n}{|x-y|^{n-2}}$;
(2) $|G(x,y)|\leq\frac{2x_ny_n}{\omega_n|x-y|^n}$;
(3) $|G(x,y)|\leq\frac{Ax_ny_n}{|x-y|^{n-2}|x-y^*|^2}$.
Proof: (1) is obvious; (2) follows by the Mean Value Theorem for Derivatives; (3) can be derived from [**14**].

Throughout the paper, let A denote various positive constants independent of the variables in question.

Proof of Theorem 1

Define the measure $dm(y')$ and the kernel $K(x,y')$ by

$$dm(y') = \frac{|f(y')|}{1+|y'|^{n+m}}dy', \quad K(x,y') = P_m(x,y')(1+|y'|^{n+m}).$$

For any $\varepsilon > 0$, there exists $R_\varepsilon > 2$, such that

$$\int_{|y'|\geq R_\varepsilon} dm(y') \leq \frac{\varepsilon}{5^{n-\alpha}}.$$

For every Lebesgue measurable set $E \subset \mathbf{R}^{n-1}$, the measure $m^{(\varepsilon)}$ defined by $m^{(\varepsilon)}(E) = m(E \cap \{x' \in \mathbf{R}^{n-1} : |x'| \geq R_\varepsilon\})$ satisfies $m^{(\varepsilon)}(\mathbf{R}^{n-1}) \leq \frac{\varepsilon}{5^{n-\alpha}}$, write

$$\begin{aligned}
v_1(x) &= \int_{|x-(y',0)|\leq 3|x|} P(x,y')(1+|y'|^{n+m})dm^{(\varepsilon)}(y'), \\
v_2(x) &= \int_{|x-(y',0)|\leq 3|x|} (P_m(x,y')-P(x,y'))(1+|y'|^{n+m})dm^{(\varepsilon)}(y'), \\
v_3(x) &= \int_{|x-(y',0)|> 3|x|} K(x,y')dm^{(\varepsilon)}(y'), \\
v_4(x) &= \int_{1<|y'|<R_\varepsilon} K(x,y')dm(y'), \\
v_5(x) &= \int_{|y'|\leq 1} K(x,y')dm(y').
\end{aligned}$$

then

$$|v(x)| \leq |v_1(x)| + |v_2(x)| + |v_3(x)| + |v_4(x)| + |v_5(x)|. \tag{2.3}$$

Let $E_1(\lambda) = \{x \in \mathbf{R}^n : |x| \geq 2, \exists t > 0, m^{(\varepsilon)}(B(x,t) \cap \mathbf{R}^{n-1}) > \lambda(\frac{t}{|x|})^{n-\alpha}\}$, therefore, if $|x| \geq 2R_\varepsilon$ and $x \notin E_1(\lambda)$, then we have

$$\begin{aligned}
|v_1(x)| &\leq \int_{x_n\leq|x-(y',0)|\leq 3|x|} \frac{2x_n}{\omega_n|x-(y',0)|^n} 2|y'|^{n+m}dm^{(\varepsilon)}(y') \\
&\leq \frac{4^{n+m+1}}{\omega_n} x_n|x|^{m+n} \int_{x_n}^{3|x|} \frac{1}{t^n} dm_x^{(\varepsilon)}(t) \\
&\leq \frac{4^{n+m+1}}{\omega_n}\left(\frac{1}{3^\alpha}+\frac{n}{\alpha}\right)\lambda x_n^{1-\alpha}|x|^{m+\alpha}. \qquad (2.4)
\end{aligned}$$

where $m_x^{(\varepsilon)}(t) = \int_{|x-(y',0)|\leq t} dm^{(\varepsilon)}(y')$.

By (1) and (3) of Lemma 2, we obtain

$$\begin{aligned}|v_2(x)| &\leq \int_{x_n\leq|x-(y',0)|\leq 3|x|}\sum_{k=0}^{m-1}\frac{2x_n|x|^k}{\omega_n}C_k^{n/2}(1)\frac{2|y'|^{n+m}}{|y'|^{n+k}}dm^{(\varepsilon)}(y')\\ &\leq \frac{4^{m+1}}{\omega_n}\sum_{k=0}^{m-1}\frac{1}{4^k}C_k^{n/2}(1)\frac{1}{5^{n-\alpha}}\varepsilon x_n|x|^m\\ &\leq \frac{4^{m+1+\alpha}}{\omega_n\cdot 3^n}\varepsilon x_n|x|^m. \qquad (2.5)\end{aligned}$$

By (1) and (3) of Lemma 2, we see that([**19**])

$$\begin{aligned}|v_3(x)| &\leq \int_{|x-(y',0)|>3|x|}\sum_{k=m}^{\infty}\frac{4x_n|x|^k}{\omega_n(2|x|)^{k-m}}C_k^{n/2}(1)\,dm^{(\varepsilon)}(y')\\ &\leq \frac{2^{m+2}}{\omega_n}\frac{\varepsilon}{5^{n-\alpha}}\sum_{k=m}^{\infty}\frac{1}{2^k}C_k^{n/2}(1)\,x_n|x|^m\\ &\leq \frac{2^{m-n+2\alpha+2}}{\omega_n}\varepsilon x_n|x|^m. \qquad (2.6)\end{aligned}$$

Write

$$\begin{aligned}v_4(x) &= \int_{1<|y'|<R_\varepsilon}[P(x,y')+(P_m(x,y')-P(x,y'))](1+|y'|^{n+m})dm(y')\\ &= v_{41}(x)+v_{42}(x),\end{aligned}$$

then

$$\begin{aligned}|v_{41}(x)| &\leq \int_{1<|y'|<R_\varepsilon}\frac{2x_n}{\omega_n|x-(y',0)|^n}2|y'|^{n+m}dm(y')\\ &\leq \frac{4R_\varepsilon^{n+m}x_n}{\omega_n}\int_{1<|y'|<R_\varepsilon}\frac{1}{(\frac{|x|}{2})^n}dm(y')\\ &\leq \frac{2^{n+2}R_\varepsilon^{n+m}m(\mathbf{R}^{n-1})}{\omega_n}\frac{x_n}{|x|^n}. \qquad (2.7)\end{aligned}$$

by (1) and (3) of Lemma 2, we obtain

$$\begin{aligned}|v_{42}(x)| &\leq \int_{1<|y'|<R_\varepsilon}\sum_{k=0}^{m-1}\frac{2x_n|x|^k}{\omega_n|y'|^{n+k}}C_k^{n/2}(1)\cdot 2|y'|^{n+m}dm(y')\\ &\leq \sum_{k=0}^{m-1}\frac{4}{\omega_n}C_k^{n/2}(1)\,x_n|x|^kR_\varepsilon^{m-k}m(\mathbf{R}^{n-1})\\ &\leq \frac{2^{n+m+1}R_\varepsilon^m m(\mathbf{R}^{n-1})}{\omega_n}x_n|x|^{m-1}. \qquad (2.8)\end{aligned}$$

In case $|y'| \leq 1$, note that

$$K(x,y') = P_m(x,y')(1+|y'|^{n+m}) \leq \frac{4x_n}{\omega_n|x-(y',0)|^n},$$

so that

$$|v_5(x)| \leq \int_{|y'|\leq 1} \frac{4x_n}{\omega_n(\frac{|x|}{2})^n} dm(y') \leq \frac{2^{n+2}m(\mathbf{R}^{n-1})}{\omega_n}\frac{x_n}{|x|^n}. \tag{2.9}$$

Thus, by collecting (2.3), (2.4), (2.5), (2.6), (2.7), (2.8) and (2.9), there exists a positive constant A independent of ε, such that if $|x| \geq 2R_\varepsilon$ and $x \notin E_1(\varepsilon)$, we have

$$|v(x)| \leq A\varepsilon x_n^{1-\alpha}|x|^{m+\alpha}.$$

Let μ_ε be a measure in $\mathbf{R}^n$ defined by $\mu_\varepsilon(E) = m^{(\varepsilon)}(E \cap \mathbf{R}^{n-1})$ for every measurable set E in $\mathbf{R}^n$.Take $\varepsilon = \varepsilon_p = \frac{1}{2^{p+2}}, p = 1, 2, 3, \cdots$, then there exists a sequence $\{R_p\}$: $1 = R_0 < R_1 < R_2 < \cdots$ such that

$$\mu_{\varepsilon_p}(\mathbf{R}^n) = \int_{|y'|\geq R_p} dm(y') < \frac{\varepsilon_p}{5^{n-\alpha}}.$$

Take $\lambda = 3 \cdot 5^{n-\alpha} \cdot 2^p \mu_{\varepsilon_p}(\mathbf{R}^n)$ in Lemma 1, then there exists $x_{j,p}$ and $\rho_{j,p}$, where $R_{p-1} \leq |x_{j,p}| < R_p$, such that

$$\sum_{j=1}^{\infty} (\frac{\rho_{j,p}}{|x_{j,p}|})^{n-\alpha} \leq \frac{1}{2^p}.$$

if $R_{p-1} \leq |x| < R_p$ and $x \notin G_p = \cup_{j=1}^{\infty} B(x_{j,p}, \rho_{j,p})$, we have

$$|v(x)| \leq A\varepsilon_p x_n^{1-\alpha}|x|^{m+\alpha},$$

Thereby

$$\sum_{p=1}^{\infty}\sum_{j=1}^{\infty} (\frac{\rho_{j,p}}{|x_{j,p}|})^{n-\alpha} \leq \sum_{p=1}^{\infty} \frac{1}{2^p} = 1 < \infty.$$

Set $G = \cup_{p=1}^{\infty} G_p$, thus Theorem 1 holds.

Proof of Theorem 2

Define the measure $dn(y)$ and the kernel $L(x,y)$ by

$$dn(y) = \frac{y_n d\mu(y)}{1+|y|^{n+m}}, \quad L(x,y) = G_m(x,y)\frac{1+|y|^{n+m}}{y_n}.$$

then the function $h(x)$ can be written as

$$h(x) = \int_H L(x,y)dn(y).$$

For any $\varepsilon > 0$, there exists $R_\varepsilon > 2$, such that

$$\int_{|y|\geq R_\varepsilon} dn(y) < \frac{\varepsilon}{5^{n-\alpha}}.$$

For every Lebesgue measurable set $E \subset \mathbf{R}^n$, the measure $n^{(\varepsilon)}$ defined by $n^{(\varepsilon)}(E) = n(E \cap \{y \in H : |y| \geq R_\varepsilon\})$ satisfies $n^{(\varepsilon)}(H) \leq \frac{\varepsilon}{5^{n-\alpha}}$, write

$$\begin{aligned}
h_1(x) &= \int_{|x-y|\leq \frac{x_n}{2}} G(x,y)\frac{1+|y|^{n+m}}{y_n}dn^{(\varepsilon)}(y),\\
h_2(x) &= \int_{\frac{x_n}{2}<|x-y|\leq 3|x|} G(x,y)\frac{1+|y|^{n+m}}{y_n}dn^{(\varepsilon)}(y),\\
h_3(x) &= \int_{|x-y|\leq 3|x|} (G_m(x,y)-G(x,y))\frac{1+|y|^{n+m}}{y_n}dn^{(\varepsilon)}(y),\\
h_4(x) &= \int_{|x-y|>3|x|} L(x,y)dn^{(\varepsilon)}(y),\\
h_5(x) &= \int_{1<|y|<R_\varepsilon} L(x,y)dn(y),\\
h_6(x) &= \int_{|y|\leq 1} L(x,y)dn(y).
\end{aligned}$$

then

$$h(x) = h_1(x)+h_2(x)+h_3(x)+h_4(x)+h_5(x)+h_6(x). \tag{2.10}$$

Let $E_2(\lambda) = \{x \in \mathbf{R}^n : |x| \geq 2, \exists t > 0, n^{(\varepsilon)}(B(x,t)\cap H) > \lambda(\frac{t}{|x|})^{n-\alpha}\}$, therefore, if $|x| \geq 2R_\varepsilon$ and $x \notin E_1(\lambda)$, then we have by (1) of Lemma 3

$$\begin{aligned}
|h_1(x)| &\leq \int_{|x-y|\leq \frac{x_n}{2}} \frac{r_n}{|x-y|^{n-2}}\frac{2|y|^{n+m}}{\frac{x_n}{2}}dn^{(\varepsilon)}(y)\\
&\leq 4\times(3/2)^{n+m}r_n\frac{|x|^{n+m}}{x_n}\int_0^{\frac{x_n}{2}}\frac{1}{t^{n-2}}dn_x^{(\varepsilon)}(t)\\
&\leq 4\times(3/2)^{n+m}r_n\left[\frac{1}{2^{2-\alpha}}+\frac{n-2}{(2-\alpha)2^{2-\alpha}}\right]\lambda x_n^{1-\alpha}|x|^{m+\alpha}. \quad (2.11)
\end{aligned}$$

where $n_x^{(\varepsilon)}(t) = \int_{|x-y|\leq t} dn^{(\varepsilon)}(y)$.

By (2) of Lemma 3, we have

$$\begin{aligned}
|h_2(x)| &\leq \int_{\frac{x_n}{2}<|x-y|\leq 3|x|} \frac{2x_n y_n}{\omega_n|x-y|^n}\frac{2|y|^{n+m}}{y_n}dn^{(\varepsilon)}(y)\\
&\leq \frac{4^{n+m+1}}{\omega_n}x_n|x|^{n+m}\int_{\frac{x_n}{2}}^{3|x|}\frac{1}{t^n}dn_x^{(\varepsilon)}(t)\\
&\leq \frac{4^{n+m+1}}{\omega_n}\left(\frac{1}{3^\alpha}+\frac{n2^\alpha}{\alpha}\right)\lambda x_n^{1-\alpha}|x|^{m+\alpha}. \quad (2.12)
\end{aligned}$$

First note $C_0^\lambda(t) \equiv 1$([**7**]) , then we obtain by (1), (3) and (4) of Lemma 2 and taking $t = \frac{x\cdot y}{|x||y|}$, $t^* = \frac{x\cdot y^*}{|x||y^*|}$ in (4) of Lemma 2

$$\begin{aligned}
|h_3(x)| &\leq \int_{|x-y|\leq 3|x|} \sum_{k=1}^{m} \frac{r_n|x|^k}{|y|^{n-2+k}} 2(n-2) C_{k-1}^{n/2}(1) \frac{x_n y_n}{|x||y|} \frac{2|y|^{n+m}}{y_n} dn^{(\varepsilon)}(y) \\
&\leq \frac{4^{m+1}}{\omega_n} \sum_{k=1}^{m} \frac{1}{4^{k-1}} C_{k-1}^{n/2}(1) \frac{1}{5^{n-\alpha}} \varepsilon x_n |x|^m \\
&\leq \frac{4^{m+1+\alpha}}{\omega_n \cdot 3^n} \varepsilon x_n |x|^m. \qquad (2.13)
\end{aligned}$$

By (1), (3) and (4) of Lemma 2, we see that

$$\begin{aligned}
|h_4(x)| &\leq \int_{|x-y|>3|x|} \sum_{k=m+1}^{\infty} \frac{r_n|x|^k}{|y|^{n-2+k}} 2(n-2) C_{k-1}^{n/2}(1) \frac{x_n y_n}{|x||y|} \frac{2|y|^{n+m}}{y_n} dn^{(\varepsilon)}(y) \\
&\leq \frac{2^{m+2}}{\omega_n} \sum_{k=m+1}^{\infty} \frac{1}{2^{k-1}} C_{k-1}^{n/2}(1) \frac{1}{5^{n-\alpha}} \varepsilon x_n |x|^m \\
&\leq \frac{2^{m-n+2\alpha+2}}{\omega_n} \varepsilon x_n |x|^m. \qquad (2.14)
\end{aligned}$$

Write

$$\begin{aligned}
h_5(x) &= \int_{1<|y|<R_\varepsilon} [G(x,y) + (G_m(x,y) - G(x,y))] \frac{1+|y|^{n+m}}{y_n} dn(y) \\
&= h_{51}(x) + h_{52}(x),
\end{aligned}$$

then we obtain by (2) of Lemma 3

$$\begin{aligned}
|h_{51}(x)| &\leq \int_{1<|y|<R_\varepsilon} \frac{2x_n y_n}{\omega_n |x-y|^n} \frac{2|y|^{n+m}}{y_n} dn(y) \\
&\leq \frac{4R_\varepsilon^{n+m}}{\omega_n} x_n \int_{1<|y|<R_\varepsilon} \frac{1}{(\frac{|x|}{2})^n} dn(y) \\
&\leq \frac{2^{n+2} R_\varepsilon^{n+m} n(H)}{\omega_n} \frac{x_n}{|x|^n}. \qquad (2.15)
\end{aligned}$$

by (1), (3) and (4) of Lemma 2, we obtain

$$\begin{aligned}
|h_{52}(x)| &\leq \int_{1<|y|<R_\varepsilon} \sum_{k=1}^{m} \frac{r_n|x|^k}{|y|^{n-2+k}} 2(n-2) C_{k-1}^{n/2}(1) \frac{x_n y_n}{|x||y|} \frac{2|y|^{n+m}}{y_n} dn(y) \\
&\leq \sum_{k=1}^{m} \frac{4}{\omega_n} C_{k-1}^{n/2}(1)\, x_n |x|^{k-1} R_\varepsilon^{m-k+1} n(H) \\
&\leq \frac{2^{n+m+1} R_\varepsilon^m n(H)}{\omega_n} x_n |x|^{m-1}. \qquad (2.16)
\end{aligned}$$

In case $|y| \leq 1$, by (2) of Lemma 3, we have

$$|L(x,y)| \leq \frac{2x_n y_n}{\omega_n |x-y|^n}\frac{2}{y_n} = \frac{4x_n}{\omega_n |x-y|^n},$$

so that

$$|h_6(x)| \leq \int_{|y|\leq 1} \frac{4x_n}{\omega_n (\frac{|x|}{2})^n} dn(y) \leq \frac{2^{n+2} n(H)}{\omega_n}\frac{x_n}{|x|^n}. \tag{2.17}$$

Thus, by collecting (2.10), (2.11), (2.12), (2.13), (2.14), (2.15), (2.16) and (2.17), there exists a positive constant A independent of ε, such that if $|x| \geq 2R_\varepsilon$ and $x \notin E_2(\varepsilon)$, we have

$$|h(x)| \leq A\varepsilon x_n^{1-\alpha}|x|^{m+\alpha}.$$

Similarly, if $x \notin G$, we have

$$h(x) = o(x_n^{1-\alpha}|x|^{m+\alpha}) \quad \text{as } |x| \to \infty. \tag{2.18}$$

by (1.11) and (2.18), we obtain

$$u(x) = v(x) + h(x) = o(x_n^{1-\alpha}|x|^{m+\alpha}) \quad \text{as } |x| \to \infty$$

hold in $H - G$, thus we complete the proof of Theorem 2.

Proof of Theorem 3 and 4

We prove only Theorem 4, the proof of Theorem 3 is similar. By (2.13), (2.14), (2.15), (2.16) and (2.17) we have

$$\lim_{|x|\to\infty, x\in H} \frac{h_3(x)+h_4(x)+h_5(x)+h_6(x)}{|x|^{m+1}} = 0. \tag{2.19}$$

In view of (1.12), we can find a sequence $\{a_i\}$ of positive numbers such that $\lim_{i\to\infty} a_i = \infty$ and

$$\sum_{i=1}^{\infty} a_i \int_{F_i} \frac{y_n}{|y|^{n+m}} d\mu(y) < \infty.$$

Consider the sets

$$E_i = \{x \in H : 2^i \leq |x| < 2^{i+1}, |h_1(x)+h_2(x)| \geq a_i^{-1} 2^{im}|x|\}$$

for $i = 1, 2, \cdots$. If $x \in E_i$, then we obtain by (3) of Lemma 3

$$a_i^{-1} \leq 2^{-im}|x|^{-1}|h_1(x)+h_2(x)| \leq A 2^{-i(m+1)} \int_{F_i} \frac{y_n}{|x-y|^{n-1}} d\mu(y)$$

so that it follows from the definition of $C(E_i; F_i)$ that

$$C(E_i; F_i) \leq A a_i 2^{-i(m+1)} \int_{F_i} y_n d\mu(y) \leq A a_i 2^{i(n-1)} \int_{F_i} \frac{y_n}{|y|^{n+m}} d\mu(y)$$

Define $E = \bigcup_{i=1}^{\infty} E_i$, then

$$\sum_{i=1}^{\infty} 2^{-i(n-1)} C(E_i; F_i) < \infty.$$

Clearly,

$$\lim_{|x|\to\infty, x\in H-E} \frac{h_1(x)+h_2(x)}{|x|^{m+1}} = 0. \tag{2.20}$$

Thus, by collecting (2.19) and (2.20), the proof of Theorem 4 is completed.

References

[1] Armitage, D. H., *Representations of harmonic functions in half-spaces.* Proc. London Math. Soc., (3) 38 (1979), no. 1, 53–71.

[2] Armitage, D. H., *Representations of superharmonic functions in half-spaces.* J. London Math. Soc. (2) 18 (1978), no. 2, 273–286.

[3] David Siegel and Erik Talvila, *Sharp growth estimates for modified Poisson integrals in a half space.* Potential Analysis, 15(2001), 333-360, MR 2002h:31008, Zbl 0987.31003.

[4] Deng Guantie, *Integral representations of harmonic functions in half spaces.* Bull.Sci.Math. 131(2007) 53-59.

[5] D.Siegel and E.Talvila, *Uniqueness for the n-dimensional half space Dirichlet problem.* Pacific J.Math, 175(1996), 571-587, MR 98a:35020, Zbl 0865.35038.

[6] E.M.Stein, *Singular integrals and differentiability properties of functions.* Princeton Univ. Press, Princeton, NJ, 1979.

[7] E.M.Stein and G.Weiss, *Introduction to Fourier Analysis on Euclidean Space.* Princeton Univ. Press, Princeton, NJ, 1971.

[8] Essén, M.; Jackson, H. L., *On the covering properties of certain exceptional sets in a half-space.* Hiroshima Math. J. 10 (1980), no. 2, 233–262.

[9] Essén, M.; Jackson, H. L.; Rippon, P. J., *On a-minimally thin sets in a half-space in $R^p, p \geq 2$.* Mathematical structure—computational mathematics—mathematical modelling, 2, 158–164, Publ. House Bulgar. Acad. Sci., Sofia, 1984.

[10] Essén, M.; Jackson, H. L.; Rippon, P. J., *On minimally thin and rarefied sets in $R^p, p \geq 2$.* Hiroshima Math. J. 15 (1985), no. 2, 393–410.

[11] Gardiner, S. J., *Representation and growth of subharmonic functions in half-spaces.* Proc. London Math. Soc. (3) 48 (1984), no. 2, 300–318.

[12] G.Szegö, *Orhhogonal Polynomials.* American Mathematical Society Colloquium Publications, Vol.23, American Mathematical Society, Providence,1975, MR 51 ♯8724, Zbl 0100.28405.

[13] H. Groemer, *Geometric Applications of Fourier series and spherical Harmonics.* Encyclopedia of Mathematics and its applications; v.61, Cambridge University Press, Cambridge, England,1996.

[14] Hiroaki Aikawwa; Matts Essén, *Potential Theory-Selected Topics.* Springer-Verlag, Berlin · Heidelberg · New York, 1996.

[15] H.Yoshida, *A type of uniqueness for the Dirichlet problem on a half-space with continuous data.* Pacific J.Math, 172(1996), 591-609, MR 97g:31007, Zbl 0853.31004.

[16] Lars Hörmander, *Notions of Convexity.* Birkhäuser, Boston · Basel · Berlin, 1994.

[17] Sheldon Axler, Paul Bourdon and Wade Ramey, *Harmonic Function Theory.* Second Edition. Springer-Verlag, New York, 1992.

[18] T.M.Flett, *On the rate of growth of mean values of holomorphic and harmonic functions.* Proc. London Math. Soc., 20(3) (1970), 749-768, MR 42 ♯3286, Zbl 0211.39203.

[19] W.K.Hayman and P.B.Kennedy, *Subharmonic Functions.* Vol. 1, London Math. Soc. Monographs, 9, Academic Press, London-New York, 1976, MR 57 ♯665, Zbl 0419.31001.

[20] Yoshihiro Mizuta and Testsu Shimomura, *Growth properties for modified Poisson integrals in a half space.* Pacific J.Math, Vol.212, No.2, 2003.

[1] Sch. Math. Sci. & Lab. Math. Com. Sys., Beijing Normal University, 100875 Beijing, The People's Republic of China

[2] Department of Public Basic Courses, Beijing Institute of Fashion and Technology, 100029 Beijing, The People's Republic of China
E-mail address: `denggt@bnu.edu.cn`